

# A note on Hermitian forms and Maschke's Theorem\*

M. Boratynski

Dipartimento di Matematica

via Orabona 4

70125 Bari, Italy

boratyn@dm.uniba.it

## Abstract

The aim of this note is to show that the "usual" proof of the Maschke Theorem in the case of real and complex field can not be generalized at least to the case of a field of rational functions on a non-singular curve.

## Introduction

Let a finite group  $G$  act linearly on a finite-dimensional vector space  $V$  over a field  $k$ . It is well known (Maschke's Theorem) that in case the characteristic of  $k$  does not divide the order of  $G$  the representation  $V$  is completely reducible. If  $k$  is a real or a complex field  $V$  admits a  $G$ -invariant hermitian form  $(\ , \ ) : V \times V \rightarrow k$  which is positively defined. Then for any  $G$ -invariant subspace  $W$  of  $V$  we obtain  $V = W \oplus W^\perp$  where  $W^\perp$  denotes the orthogonal complement of  $W$  with respect to  $(\ , \ )$ . The decomposition  $V = W \oplus W^\perp$  holds since the restriction of  $(\ , \ )$  to  $W$  is nondegenerate. Moreover  $W^\perp$  is  $G$ -invariant because  $(\ , \ ) : V \times V \rightarrow k$  is  $G$ -invariant.

The natural question arises whether every finite-dimensional  $G$ -module admits a  $G$ -invariant hermitian form such that its restriction to any (proper)

---

\*Research supported by MURST 60%

$G$ -invariant submodule of  $V$  is nondegenerate in case the characteristic of  $k$  does not divide the order of  $G$ . If this were true we would obtain an alternative proof of the Maschke Theorem. The aim of this note is to show that the answer is negative even when the action of  $G$  is trivial.

In the sequel  $k$  will denote a field with a fixed involution ie. an automorphism  $\lambda \rightarrow \bar{\lambda}$  of order  $\leq 2$ . We shall always assume that the involution is not an identity or  $\text{ch } k \neq 2$  otherwise.  $V$  will denote a finite-dimensional vector space over  $k$ .

**Definition 1** A map  $(\ , \ ) : V \times V \rightarrow k$  is called a hermitian form if it is linear with respect to the first variable and moreover  $\overline{(v, w)} = (w, v)$  for all  $v, w \in V$

**Remark 1** Of course we immediately obtain  $(v, w_1 + w_2) = (v, w_1) + (v, w_2)$  and  $(v, \lambda w) = \bar{\lambda}(v, w)$  for  $v, w, w_1, w_2 \in V$  and  $\lambda \in k$ . In case the involution is an identity automorphism then obviously the notions of hermitian and symmetric forms coincide.

The following Lemma and Proposition are of course well known; the proofs are enclosed only for the completeness sake.

**Lemma** Let  $(\ , \ ) : V \times V \rightarrow k$  be a non-zero hermitian form. Then there exists  $v \in V$  such that  $(v, v) \neq 0$ .

*Proof:* Suppose  $(v, v) = 0$  for all  $v \in V$ . Then

$$0 = (v + w, v + w) = (v, w) + (w, v) \quad (1)$$

for all  $v, w \in V$ . If the involution of  $k$  is an identity we obtain  $2(v, w) = 0$ .

So  $(v, w) = 0$  for all  $v, w \in V$  since  $\text{ch } k \neq 2$  which is impossible.

Let  $\lambda \in V$  such that  $\lambda \neq \bar{\lambda}$  in case the involution is not an identity. Then

$$0 = (\lambda v, w) + (w, \lambda v) = \lambda(v, w) + \bar{\lambda}(w, v) \quad (2)$$

It follows from (1) and (2) that

$$(\lambda - \bar{\lambda})(v, w) = \lambda(v, w) + \bar{\lambda}(w, v) = 0$$

So  $(v, w) = 0$  for all  $v, w \in V$  which contradicts our hypothesis. Thus in both cases there exists  $v \in V$  such that  $(v, v) \neq 0$ .

**Proposition 1** Let  $(\ , \ ) : V \times V \rightarrow k$  be a hermitian form. Then there exist  $e_1, e_2, \dots, e_n \in V$  which form a basis of  $V$  and such that  $(e_i, e_j) = 0$  if  $i \neq j$ .  
*Proof:* If  $\dim V = 1$  then every  $v \in V$  such that  $v \neq 0$  will do. Let  $\dim V = n > 1$ . We can assume that  $(\ , \ ) : V \times V \rightarrow k$  is non-zero. By the Lemma there exists  $e \in V$  such that  $(e, e) \neq 0$ . Let  $\langle e \rangle$  denote the subspace generated by  $e$ . Put  $\langle e \rangle^\perp = \{v \in V \mid (v, e) = 0\}$  which is a subspace of  $V$ . Let  $v \in V$ . We obtain

$$(v - ((v, e)/(e, e))e, e) = (v, e) - ((v, e)/(e, e))(e, e) = (v, e) - (v, e) = 0$$

So

$$v = ((v, e)/(e, e))e + [v - ((v, e)/(e, e))e]$$

with

$$((v, e)/(e, e))e \in \langle e \rangle \text{ and } [v - ((v, e)/(e, e))e] \in \langle e \rangle^\perp.$$

Thus  $V = \langle e \rangle \oplus \langle e \rangle^\perp$  because  $\langle e \rangle \cap \langle e \rangle^\perp = (0)$ . By the inductive hypothesis applied to  $\langle e \rangle^\perp$  which is of dimension  $n - 1$  there exist  $e_2, e_3, \dots, e_n \in \langle e \rangle^\perp$  which form a basis of  $\langle e \rangle^\perp$  and such that  $(e_i, e_j) = 0$  for  $i, j = 2, 3, \dots, n$  and  $i \neq j$ . Put  $e_1 = e$ . Then  $e_1, e_2, \dots, e_n$  satisfy the required conditions.

Put  $k_0 = \{\lambda \in k \mid \bar{\lambda} = \lambda\}$ . Then  $k_0$  is a subfield of  $k$  such that  $\dim k_{k_0} \leq 2$ . For any  $\lambda \in k$  define  $\varphi(\lambda) = \lambda\bar{\lambda}$ . Then  $\varphi: k^* \rightarrow k_0^*$  and is a homomorphism where for any field  $K$ ,  $K^*$  denotes the multiplicative group of its non-zero elements. In case the involution of  $k$  is not an identity  $\varphi(\lambda) = \prod_{\theta \in G(k/k_0)} \theta(\lambda)$  where  $G(k/k_0)$  is the Galois group of the extension  $k/k_0$ . So  $\varphi$  coincides with the "usual" norm homomorphism  $N_{k/k_0}: k^* \rightarrow k_0^*$ .

**Definition 2** Let  $(\ , \ ) : V \times V \rightarrow k$  be a hermitian form. Then  $v \in V$  is called isotropic if  $(v, v) = 0$

**Proposition 2** Let  $(\ , \ ) : V \times V \rightarrow k$  be a hermitian form. If  $\varphi: k^* \rightarrow k_0^*$  is an epimorphism and  $\dim V \geq 2$  then  $V$  contains a non-zero isotropic element.

*Proof:* Let  $e_1, e_2, \dots, e_n \in V$  form a basis of  $V$  and are such that  $(e_i, e_j) = 0$  if  $i \neq j$  (Proposition 1). Put  $\lambda_i = (e_i, e_i)$  for  $i = 1, 2, \dots, n$ . We have  $(e_i, e_i) = \overline{(e_i, e_i)}$ . So  $\lambda_i \in k_0$  for  $i = 1, 2, \dots, n$ . If  $\lambda_i = 0$  for some  $i$  then the corresponding  $e_i$  is isotropic. So we can assume that  $\lambda_i \neq 0$  for  $i = 1, 2, \dots, n$ . Let us consider  $v = xe_i + e_j$  where  $x \in k$  such that  $\varphi(x) = -\lambda_j/\lambda_i$  and  $i \neq j$ . Then  $v \in V$  and  $v \neq 0$ . Moreover

$$(v, v) = (xe_i + e_j, xe_i + e_j) = x\bar{x}\lambda_i + \lambda_j = \varphi(x)\lambda_i + \lambda_j = (-\lambda_j/\lambda_i)\lambda_i + \lambda_j = 0$$

**Definition 3** A field  $K$  is called  $C_1$  ([1]) if every equation  $f(x_1, x_2, \dots, x_n) = 0$  where  $f$  is a homogenous polynomial of degree  $d < n$  with coefficients in  $K$ , admits a non-trivial solution in  $K^n$ .

**Theorem** Let  $(\ , \ ) : V \times V \rightarrow k$  be a hermitian form. If  $k_0$  is  $C_1$  and  $\dim V \geq 3$  then  $V$  admits a non-zero isotropic element.

*Proof:*  $N_{k/k_0} : k^* \rightarrow k_0^*$  is an epimorphism since  $k_0$  is  $C_1$  ([1]). It follows that  $\varphi : k^* \rightarrow k_0^*$  is an epimorphism in case the involution is not an identity. So we can apply Proposition 2. Suppose now that the involution is an identity and let  $e_1, e_2, \dots, e_n \in V$  form a basis of  $V$  and are such that  $(e_i, e_j) = 0$  if  $i \neq j$  (Proposition 1). The equation

$$\lambda_1 x_1^2 + \lambda_2 x_2^2 + \dots + \lambda_n x_n^2 = 0$$

has a non-trivial solution  $(x_1, x_2, \dots, x_n)$  in  $k = k_0$  where  $\lambda_i = (e_i, e_i)$  and  $n \geq 3$ . Let  $v = x_1 e_1 + x_2 e_2 + \dots + x_n e_n \in V$ . Then  $v \neq 0$  and  $(v, v) = \lambda_1 x_1^2 + \lambda_2 x_2^2 + \dots + \lambda_n x_n^2 = 0$ .

**Corollary** Let  $k$  be a field of transcendence degree one over an algebraically closed field  $F$  with an involution which is an identity on  $F$ . Suppose  $(\ , \ ) : V \times V \rightarrow k$  is a hermitian form. If  $\dim V \geq 3$  then  $V$  admits a non-trivial isotropic element.

*Proof:* If  $k$  is of transcendence degree one over  $F$  then the same holds for  $k_0$  since  $k$  is a finite extension of  $k_0$ . By the Tsen Theorem ([1])  $k_0$  is  $C_1$ , so the Theorem can be applied.

**Remark 2** Suppose  $k$  is a field with involution which satisfies the assumptions of the Corollary and let  $(\ , \ ) : V \times V \rightarrow k$  be a hermitian form on a vector space  $V$  with a trivial action of a finite group  $G$ . Then obviously  $W \subset W^\perp$  where  $W$  denotes (a  $G$ -invariant) subspace generated by a non-zero isotropic element of  $V$ . So in case  $\dim V \geq 3$  and the involution is an identity on  $F$  (a condition very natural if  $k$  is considered a field of rational functions on a non-singular curve over  $F$ ) the proof of Maschke's Theorem in case of a real or complex field can not be generalized.

## References

- [1] J. P Serre, *Corps Locaux*, Hermann, Paris 1968.